

INDIANA BAT MIST NETTING GUIDELINES

RATIONALE

A typical mist net survey is an attempt to determine presence or probable absence of the species, it does not provide sufficient data to determine population size or structure. Following these guidelines will standardize procedures for mist netting. It will help maximize the potential for capture of Indiana bats at a minimum acceptable level of effort. Although the capture of bats confirms their presence, failure to catch bats does not absolutely confirm their absence. Netting effort as extensive as outlined below usually is sufficient to capture Indiana bats. However, there have been instances in which additional effort was necessary to detect the presence of the species.

NETTING SEASON

May 15 - August 15

These dates define acceptable limits for documenting the presence of summer population of Indiana bats, especially maternity colonies. Several captures, including adult females and young of the year, indicate that a nursery colony is active in the area. Outside these dates, even when Indiana bats are caught, data should be carefully interpreted: If only a single bat is captured, it may be a transient or migratory individual.

EQUIPMENT

Mist nets - Use the finest, lowest visibility mesh commercially available:

1. In the past, this was 1 ply, 40 denier monofilament - denoted 40/1
2. Currently, monofilament is not available and the finest on the market is 2 ply, 50 denier nylon - denoted 50/2
3. Mesh of approximately $1 \frac{1}{2}$ ($1 \frac{1}{4}$ - $1 \frac{3}{4}$) in (~38 mm)

Hardware - No specific hardware is required. There are many suitable systems of ropes and/or poles to hold the nets. See NET PLACEMENT below for minimum net heights, habitats, and other netting requirements that affect the choice of hardware. The system of Gardner, *et al.* (1989) has met the test of time.

NET PLACEMENT

Potential travel corridors such as streams or logging trails typically are the most effective places to net. Place the nets approximately perpendicular across the corridor. Nets should fill the corridor from side to side and from stream (or ground) level up to the overhanging canopy. A typical set is seven meters high consisting of three or more nets "stacked" on top one another and up to 20 m wide. (Different width nets may be purchased and used as the situation dictates.)

Occasionally it may be desirable to net where there is no good corridor. Take caution to get the nets up into the canopy. The typical equipment described in the section above may be inadequate for these situations, requiring innovation on the part of the observers.

RECOMMENDED NET SITE SPACING:

Stream corridors - one net site per km of stream.

Non-corridor land tracts - two net sites per square km of forested habitat
(= 1 net site for every 123 acres of forested habitat)

MINIMUM LEVEL OF EFFORT

Netting at each site should consist of:

At least four net-nights (unless bats are caught sooner) (one net set up for one night = one net-night)

A minimum of two net locations at each site (at least 30m apart, especially in linear habitat such as a stream corridor)

A minimum of two nights of netting

Sample Period: begin at sunset; net for at least 5 hr

Each net should be checked approximately every 20 min

No disturbance near the nets, other than to check nets and remove bats

WEATHER CONDITIONS

Severe weather adversely affects capture of bats. If Indiana bats are caught during weather extremes, it is probably because they are at the site and active despite inclement weather. On the other hand, if bats are not caught, it may be that there are bats at the site but they may be inactive due to the weather. Negative results combined with any of the following weather conditions throughout all or most of a sampling period are likely to require additional netting:

- Precipitation
- Temperatures below 10°C
- Strong winds (Use good judgement: moving nets are more likely to be detected by bats.)

MOONLIGHT

There is some evidence that small myotis bats avoid brightly lit areas, perhaps as predator avoidance. It is typically best to set nets under the canopy where they are out of the moon light, particularly when the moon is ½-full or greater.